

# Iso-Representation of the Deuteron Spin and Magnetic Moment via Bohm's Hidden Variables

Ruggero Maria Santilli

The Institute for Basic Research, 35246 U. S. 19N, Suite 215, Palm Harbor, FL 34684, USA.  
E-mail: research@i-b-r.org

In this paper, we review and upgrade the iso-representation of the spin  $1/2$  of nucleons according to the isotopic branch of hadronic mechanics, known as *hadronic spin*, which is characterized by an isotopy of Pauli's matrices with an explicit and concrete realization of Bohm's *hidden variable*  $\lambda$  and show, apparently for the first time, that it allows a consistent and time invariant representation of the spin  $J_D = 1$  of the Deuteron in its true ground state, that with null angular contributions  $L_D = 0$ . We then show, also apparently for the first time, that the indicated hadronic spin allows a numerically exact and time invariant representation of the magnetic moment of the Deuteron with the numeric value  $\lambda = 2.65557$ .

## 1 The Einstein-Podolsky-Rosen argument

In the preceding paper [1], we have outlined the axiom-preserving completion of 20th century applied mathematics into *iso-mathematics*, (see [2] for an extended presentation and [5–7] for independent studies), and the related *iso-mechanical branch of hadronic mechanics* (see [3] for a detailed treatment, [8–10] for independent studies and [11–13] for recent reviews) which isotopic methods have been used for the verification in [14, 15] of the 1935 historical argument by A. Einstein, B. Podolsky and N. Rosen that *Quantum mechanics is not a complete theory* [16] (see [17] for the proceedings of the 2020 *Teleconference in the EPR Argument*, and its overviews [18, 19]).

Via the use of said isotopic methods, [1] achieved, apparently for the first time, a non-relativistic and relativistic representation of *all* characteristics of the muons (including their recently measured anomalous magnetic moment) as an extended and naturally unstable hadronic bound state of electrons and positrons produced free in the spontaneous decay with the lowest mode.

In the subsequent paper [20], we showed that said isotopic methods confirm the 1983 experimentally unresolved deviations [21] from the conventional formulation of time dilation for composite particles such as the muons, in favor of its axiom-preserving isotopic completion. We indicated in [20] that said deviations are due to incompatibility of the conventional time dilation with the time-irreversible character of the muon decay voiced since 1967 by R. M. Santilli [22] (see the 1995 full treatment [3]) and independently voiced in 1968 by D. I. Blokhintsev [23] for the incompatibility of the conventional time dilation with internal non-local effects of composite particles.

In this paper, we review and upgrade the notion of *hadronic spin* first introduced in [3, Section 6.8, page 250] and then used for verification [14] of the EPR argument [16] as well as in other applications [4]. The new notion of hadronic spin is

then used for the characterization of the spin  $1/2$  of the nucleons, and realized via an isotopy of Pauli's matrices with an explicit and concrete realization of Bohm's *hidden variable*  $\lambda$  [43]. We then show, apparently for the first time, that said hadronic spin allows the first known exact and time-invariant representation of the spin  $S_D = 1$  of the deuteron in the true ground state, that with null contributions from angular momenta  $L_D = 0$ .

We then show, also apparently for the first time, that said hadronic spin allows a numerically exact and time-invariant representation of the magnetic moment of the Deuteron with  $\lambda = 2.65557$ .

A technical understanding of this paper requires a technical knowledge at least of [2, 3]. A preliminary understanding of this paper requires a knowledge of reviews [11–13].

## 2 Iso-representation of the Deuteron spin

As it is well known, the quantum mechanical spin  $1/2$  of nucleons is characterized by the fundamental irreducible representation of the special unitary Lie algebra  $SU(2)$  which is notoriously given by the celebrated *Pauli matrices*

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad (1)$$

(where  $\sigma_3$  is set hereon along the spin direction) with commutation rules

$$[\sigma_i, \sigma_j] = \sigma_i \sigma_j - \sigma_j \sigma_i = i2\epsilon_{ijk}\sigma_k. \quad (2)$$

The value  $S = 1/2$  of the nucleon spin is characterized by the eigenvalue equations on a Hilbert space  $\mathcal{H}$  over the field of complex numbers  $\mathcal{C}$  with basis  $|b\rangle$

$$\begin{aligned} S_k &= \frac{1}{2} \sigma_k, \\ \sigma_3 |b\rangle &= \pm |b\rangle, \\ \sigma^2 |b\rangle &= (\sigma_1 \sigma_1 + \sigma_2 \sigma_2 + \sigma_3 \sigma_3) |b\rangle = 3 |b\rangle. \end{aligned} \quad (3)$$

A serious insufficiency of quantum mechanics in nuclear physics, which is fully supportive of the EPR argument [16], is that the representation of the spin 1/2 of nucleons via Pauli matrices does not allow a representation of the Deuteron spin  $S_D = 1$  under the conditions of its experimental detection, that is, in its ground state with null orbital contributions  $L_D = 0$ . In fact, the sole possible *stable* bound state between a proton and a neutron permitted by quantum mechanics (qm) is the singlet

$$D = (n_{\uparrow}, n_{\downarrow})_{qm}, \quad (4)$$

for which the total spin is null,  $J_D = 0$ . In an attempt of resolving this insufficiency while preserving quantum mechanics, nuclear physicists have assumed for about one century that the Deuteron is a bound state of a proton and a neutron in *excited orbits* such that  $L_D = 1$  (see e.g. [25]).

When at Harvard University with DOE support, R. M. Santilli noted that the most effective way of resolving the above and other insufficiencies of quantum mechanics (see next section) is to *exit from its class of unitary equivalence*. Therefore, Santilli proposed in two 1978 memoirs [26, 27] and in two Springer Verlag monographs [28, 29], the EPR generalization / completion of quantum mechanics into a new discipline which he called *hadronic mechanics* (see the Abstract and [27, pages 684,749,777] and [29, page 112]).

Hadronic mechanics was conceived to be an *axiom-preserving, thus isotopic non-unitary image of quantum mechanics for the representation of the dimension, shape and density of hadrons in interior conditions with ensuing potential as well as non-potential interactions due to mutual penetration*.

The proposal voiced in [26]–[29] suggested the construction of the *time irreversible completion of quantum mechanics into hadronic mechanics with the basic time evolution* (see [27, (4.15.24), page 742], [29, (19), page 153] and [3, (4.3.1), page 154])

$$\begin{aligned} i \frac{dA}{dt} &= (A, H) = ARH - HSA = \\ &= (ATH - HTA) + (AJH + HJA), \\ A(t) &= e^{HSit} A(0) e^{-iRH}, \\ R &= T + J, \quad s = -T + J, \end{aligned} \quad (5)$$

which is called Lie-admissible / Jordan-admissible since the bracket  $(A, H)$  clearly contains a Lie algebra  $(ATH - HTA)$  and a Jordan algebra  $(AJH + HJA)$  content.

By recalling that quantum mechanics can only represent systems whose time reversal images verify causality laws (because Heisenberg's equation is invariant under anti-Hermiticity), the aim of Santilli's proposal (stemming from his DOE support) was to achieve a consistent treatment of systems whose time reversal image violate causality, which is the case for all energy-releasing processes, with particular reference to nuclear fusions and fossil fuel combustion.

In this paper, we study *stable nuclei* that, as such, are time-reversal invariant. Consequently, our study requires the *Lie-isotopic branch of hadronic mechanics*, called for brevity *iso-mechanics*, which is based on the completion of the quantum mechanical enveloping associative algebra of Hermitean operators  $A, B, \dots$  on  $\mathcal{H}$  over  $\mathcal{C}$  with product  $A \times B = AB$  and multiplicative unit  $I$  into the new product (first introduced in [26, (3.710), page 352] and [29, (5), page 71])

$$A \star B = A\hat{T}B, \quad \hat{T} > 0, \quad (6)$$

called *iso-product* because associativity-preserving, the positive-definite quantity  $\hat{T}$  being called the *isotopic element* and new compatible multiplicative unit

$$\hat{I} = 1/\hat{T} > 0, \quad \hat{I} \star A = A \star \hat{I} \equiv A \forall A \in \mathcal{H}, \quad (7)$$

called *iso-unit* with ensuing basis time evolution first introduced in [27, (4.15.59), page 752] (see also [29, (18), page 163, Vol. II] and [3, (3.1.6), page 81])

$$i \frac{dA}{dt} = [A, H]^{\dagger} = A\hat{T}H - H\hat{T}A, \quad (8)$$

$$A(t) = e^{H\hat{T}it} A(0) e^{-i\hat{T}H} = W(t) A(0) W(t)^{\dagger}, \quad (9)$$

$$WW^{\dagger} \neq I,$$

which is called Lie-isotopic because of the clear verification of the Lie algebra axioms by the new brackets  $[A, H]^{\star}$ , although in a generalized form.

Following the identification of the basic structure (6) to (8), Santilli constructed in the 1983 monograph [29] the systematic isotopies in the 1983 volume [29] of the various branches of Lie's theory (universal enveloping associative algebra, Lie's theorems, Lie's transformation groups, etc.), resulting in a theory nowadays known as the *Lie-Santilli iso-theory* [5] (see also [30, 31]).

Santilli then constructed the isotopies of all known space-time symmetries [32]–[42]. In particular, systematic studies were conducted on the construction, classification and verification isotopies of the SU(2)-spin symmetry which can be found in [3, Chapter 6, page 209 on], in papers [33]–[37] with a summary in Section 3 of [12].

The *hadronic spin* (first introduced in [3, Section 6.8]) is the characterization of the spin of hadrons under strong interactions via the iso-irreducible, iso-unitary, iso-representations of the Lie-Santilli iso-symmetry  $\widehat{\text{SU}}(2)$ .

The simplest possible case of spin 1/2 of the nucleons can be outlined as following: all mathematical and physical aspects of the (regular [31]) isotopic branch of hadronic mechanics can be uniquely and unambiguously constructed via a simple, positive-definite *non-unitary transformation* set equal to the iso-unit of the new theory

$$UU^{\dagger} = \hat{I} > 0 \neq I, \quad \hat{T} = 1/\hat{I} = (UU^{\dagger})^{-1} > 0, \quad (10)$$

provided said non-unitary transformation is applied to the *totality* of the quantum mechanical, mathematical and physical quantities and their operations *with no exception known to the author*, to prevent insidious inconsistencies in mixing mathematics and iso-mathematics that generally remain undetected by non-experts in the field.

The indicated correct use of the above procedure permits the map of all quantum mechanical quantities, including unit, product, Lie algebras, etc., into their hadronic formulations that are generally denoted with a “hat”

$$\begin{aligned} I &\rightarrow UIU^\dagger = \hat{I}, \\ AB &\rightarrow U(AB)U^\dagger = \\ &= (UAU^\dagger)(UU^\dagger)^{-1}(UBU^\dagger) = \hat{A} \star \hat{B}, \quad (11) \\ AB - BA &= [A, B] \rightarrow U(AB - BA)U^\dagger = \\ &= \hat{A} \star \hat{B} - \hat{B} \star \hat{A} = [\hat{A}, \hat{B}]^*, \text{ etc.} \end{aligned}$$

The hadronic spin 1/2 for nuclear constituents is given by the iso-fundamental, iso-unitary, iso-irreducible iso-representation of the Lie-Santilli iso-algebra  $\widehat{SU}(2)$  under the condition of *iso-unimodularity*

$$\text{Det } \hat{I} = 1. \quad (12)$$

The above condition allowed Santilli to characterize the basic iso-unit of iso-mechanics in terms of Bohm's *hidden variable*  $\lambda$  [43] which was presented for the first time in [3, (6.8.19), page 248], according to the rules

$$\begin{aligned} \text{Det } \hat{I} &= \text{Det} [(UU^\dagger)] = \text{Det} [\text{Diag} (g_{11}, g_{22})] = 1, \\ g_{11} &= g_{22}^{-1} = \lambda \geq 0, \end{aligned} \quad (13)$$

yielding the *iso-Pauli matrices* first proposed in [3, (6.8.20), page 248]

$$\begin{aligned} \hat{\sigma}_k &= U\sigma_k U^\dagger, \\ UU^\dagger &= \hat{I} = \text{Diag} (\lambda^{-1}, \lambda), \quad \hat{T} = \text{Diag} (\lambda, \lambda^{-1}), \\ \hat{\sigma}_1 &= \begin{pmatrix} 0 & \lambda \\ \lambda^{-1} & 0 \end{pmatrix}, \quad \hat{\sigma}_2 = \begin{pmatrix} 0 & -i\lambda \\ i\lambda^{-1} & 0 \end{pmatrix}, \\ \hat{\sigma}_3 &= \begin{pmatrix} \lambda^{-1} & 0 \\ 0 & -\lambda \end{pmatrix}, \end{aligned} \quad (14)$$

and then used in [14] for the verification of the EPR argument thanks to the evident inapplicability of Bell's theorem [44] due to the non-unitary structure of the theory.

It is easy to see that the iso-Pauli matrices verify the Lie-Santilli iso-commutation rules

$$\begin{aligned} [\hat{\sigma}_i, \hat{\sigma}_j]^* &= \hat{\sigma}_i \star \hat{\sigma}_j - \hat{\sigma}_j \star \hat{\sigma}_i = \\ &= \hat{\sigma}_i \hat{T} \hat{\sigma}_j - \hat{\sigma}_j \hat{T} \hat{\sigma}_i = i2\epsilon_{ijk} \hat{\sigma}_k, \end{aligned} \quad (15)$$

showing the clear iso-morphism  $\widehat{SU}(2) \approx SU(2)$ .

The representation of the spin 1/2 of nucleons despite its generalized structure is given by the iso-eigenvalues on an iso-state  $|\hat{b}\rangle$  of the *Hilbert-Myung-Santilli iso-space*  $\hat{\mathcal{H}}$  [45] over the iso-field of iso-complex iso-numbers  $\hat{C}$  [46]

$$\begin{aligned} \hat{S}_k &= \hat{I} \star \hat{\sigma}_k = \frac{1}{2} \hat{\sigma}_k, \\ \hat{\sigma}_3 \star |\hat{b}\rangle &= \hat{\sigma}_3 \hat{T} |\hat{b}\rangle = \pm |\hat{b}\rangle, \\ \hat{\sigma}^2 \star |\hat{b}\rangle &= (\hat{\sigma}_1 \hat{T} \hat{\sigma}_1 + \hat{\sigma}_2 \hat{T} \hat{\sigma}_2 + \hat{\sigma}_3 \hat{T} \hat{\sigma}_3) \hat{T} |\hat{b}\rangle = 3 |\hat{b}\rangle. \end{aligned} \quad (16)$$

As it is well known, *non-unitary theories violate causality*, and that is the case for the hadronic spin when considered in its projection on a conventional Hilbert space  $\mathcal{H}$  over a conventional field  $C$ . Additionally, non-unitary transforms generally change the numeric value of the isotopic element which represent physical, measurable quantities (see next section). These and other problems are resolved by the reformulation of non-unitary time evolution (9) into the *iso-unitary iso-transformations* [47]

$$\begin{aligned} WW^\dagger &= \hat{I}, \quad W = \hat{W} \hat{T}^{1/2}, \\ WW^\dagger &= \hat{W} \star \hat{W}^\dagger = \hat{W}^\dagger \star \hat{W} = \hat{I}, \end{aligned} \quad (17)$$

under which reformulation the iso-unit, iso-product, Lie-Santilli iso-algebras, etc., are invariant,

$$\hat{I} \rightarrow \hat{W} \star \hat{I} \star \hat{W}^\dagger = \hat{I}' \equiv \hat{I}, \quad (18)$$

$$\begin{aligned} \hat{A} \star \hat{B} &\rightarrow \hat{W} \star (\hat{A} \star \hat{B}) \star \hat{W}^\dagger = \\ &= \hat{A}' \star \hat{B}' = \hat{A}' \hat{T}' \hat{B}', \quad \hat{T}' \equiv \hat{T}, \\ \hat{A}' &= \hat{W} \star \hat{A} \star \hat{W}^\dagger, \quad \hat{B}' = \hat{W} \star \hat{B} \star \hat{W}^\dagger, \\ \hat{T}' &= (\hat{W}^\dagger \star \hat{W})^{-1}. \end{aligned} \quad (19)$$

It should be noted that, by no means, hadronic spin solely characterizes the spin 1/2 because it was conceived [26, 27] for the characterization of the most general possible notion of spin for an extended particle such as a hadron in the core of a star with ensuing *non-local contributions from the star environment* (see [3, 14], [34]–[37]) according to the *de Broglie-Bohm non-local theory* [48]. The notion of hadronic spin was then specialized to the spin of nucleons because of clear experimental evidence, rather than popular views in nuclear physics, establishing its value 1/2.

The iso-representation of the Deuteron spin  $J_D = 1$  in its true bound state with  $L_D = 0$  via the hadronic spin is elementary. To see it, let us call for clarity *iso-protons, iso-neutron, iso-nucleons, iso-Deuteron and iso-Helium* (with corresponding symbols  $\hat{p}, \hat{n}, \hat{N}, \hat{D}, \hat{He}$ ), the particles and nuclei characterized by the hadronic spin. With reference to [3, Section 2.11, page 265 on] on the addition of hadronic spins, the most stable hadronic bound state of the iso-Deuteron as a

hadronic bound state of an iso-proton and an iso-neutron is given by the *axial triplet state*. The axial triplet coupling first identified in the new chemical species of magnecules (see [49, Chapter 8, page 303 on] and [50, 51]) and then used for the new Intermediate Controlled Nuclear Fusion [52–54] with iso-representation (Fig. 1)

$$\hat{D} = \begin{pmatrix} \hat{p}_\uparrow \\ \star \\ \hat{n}_\uparrow \end{pmatrix}. \quad (20)$$

### 3 Iso-representation of the Deuteron magnetic moment

Another serious limitation of quantum mechanics in nuclear physics has been the inability, in about one century of studies, to achieve an exact representation of nuclear magnetic moments via the tabulated values for the magnetic moments of the proton and of the neutron in vacuum [55]

$$\mu_p = +2.79285 \mu_N, \quad \mu_n = -1.91304 \mu_N, \quad (21)$$

where  $\mu_N$  represents the *nuclear magneton*.

As an example, the magnetic moment predicted by quantum mechanics (qm) from values (21) for the magnetic moment of the Deuteron is given by

$$\mu_D^{qm} = \mu_p + \mu_n = (2.79285 - 1.91304) \mu_N = 0.87981 \mu_N, \quad (22)$$

and does not represent the experimental value of the Deuteron magnetic moment

$$\mu_D^{ex} = 0.85647 \mu_N, \quad (23)$$

due to a deviation in *excess* of about 3%,

$$\mu_D^{qm} - \mu_D^{ex} = 0.02334 \mu_N \approx 2.95\% \mu_D^{ex}, \quad (24)$$

with larger deviations for heavier nuclei.

E. Fermi [56], V.F. Weisskopf [25] and other founders of nuclear physics formulated the hypothesis, hereon referred to as the *Fermi-Weisskopf hypothesis*, that in the transition from isolated particles in vacuum to members of a nuclear structure, protons and neutrons experience a deformation of their extended charge distribution with consequential change of their magnetic moments (21) while conserving their spin 1/2 (see the statement at the top of [25, page 31]).

The first numerically exact and time-invariant representation of the Deuteron magnetic moment (23) was achieved in 1994 by Santilli [57] (see also its subsequent extended study in [58]) thanks to the prior construction of the isotopic branch of hadronic mechanics for the representation of extended, thus deformable hadrons and related iso-symmetries [32]–[42] with the isotopic element

$$\hat{T} = \text{Diag} \left( \frac{1}{n_1^2}, \frac{1}{n_2^2}, \frac{1}{n_3^2}, \frac{1}{n_4^2} \right), \quad (25)$$

in which  $n_k^2$ ,  $k = 1, 2, 3$ , represent the semi-axes of the *deformable* proton and of the nucleon under strong nuclear forces and  $n_4^2$  represents their density. Under the assumption, for simplicity, that the proton and the neutron in the Deuteron structure have the same dimension, shape and density, [57] reached a numerically exact and time invariant representation of the magnetic moment of the Deuteron in [57, (3.6), page 124] with the following values of the characteristic  $n$ -quantities (that are denoted with the symbols  $b_\mu = 1/n_\mu$  in [57])

$$\begin{aligned} b_1 = \frac{1}{n_1} = b_2 = \frac{1}{n_2} = 1.0028, \\ b_3 = \frac{1}{n_3} = 1.662, \quad b_4 = \frac{1}{n_4} \end{aligned} \quad (26)$$

(whose derivation is not reviewed here for brevity), by therefore confirming the 1981 preliminary experimental verification of the Fermi-Weisskopf hypothesis via neutron interferometry [59].

In this paper, we present, apparently for the first time, a second numerically exact and time invariant representation of the magnetic moment of the Deuteron (23), with spin  $S_D = 1$  in its ground state via the representation of Santilli's iso-Pauli matrices (14) by using the Clifford's algebra representation of the conventional Pauli matrices [60]–[64], whose representation is here assumed to be known for brevity.

Note that, when formulated on their associative enveloping algebra, the iso-Pauli matrices satisfy all algebraic properties of the conventional Pauli matrices. Consequently, we can use the conventional representation in its entirety and introduce the *representation of iso-Pauli matrices (14) in terms of Clifford algebra*  $\tilde{\mathbf{G}}_3 = \tilde{\mathbf{G}}_3(\mathbf{R}^3)$  with the iso-basis

$$\tilde{\mathbf{G}}_3 : \{1, \hat{\sigma}_1, \hat{\sigma}_2, \hat{\sigma}_3, \hat{\sigma}_1\hat{\sigma}_2, \hat{\sigma}_1\hat{\sigma}_3, \hat{\sigma}_2\hat{\sigma}_3, i := \hat{\sigma}_1\hat{\sigma}_2\hat{\sigma}_3\}, \quad (27)$$

and main properties

$$\begin{aligned} \hat{\sigma}_1^2 = \hat{\sigma}_2^2 = \hat{\sigma}_3^2 = 1, \\ \hat{\sigma}_{12} = \hat{\sigma}_1\hat{\sigma}_2 = -\hat{\sigma}_{21}, \quad \hat{\sigma}_{13} = \hat{\sigma}_1\hat{\sigma}_3, \quad \hat{\sigma}_{23} = \hat{\sigma}_2\hat{\sigma}_3, \\ \hat{\sigma}_{12}^2 = \hat{\sigma}_1\hat{\sigma}_2\hat{\sigma}_1\hat{\sigma}_2 = -\hat{\sigma}_1\hat{\sigma}_2\hat{\sigma}_2\hat{\sigma}_1 = -\hat{\sigma}_1^2\hat{\sigma}_2^2 = -1. \end{aligned} \quad (28)$$

The *standard basis* of unit iso-vectors  $\{\hat{\sigma}_1, \hat{\sigma}_2, \hat{\sigma}_3\}$  define the  $x, y, z$  iso-coordinate axes, respectively. The *iso-spectral basis* is

$$\begin{pmatrix} \hat{u}_+ & \hat{\sigma}_1\hat{u}_- \\ \hat{\sigma}_1\hat{u}_+ & \hat{u}_- \end{pmatrix}, \quad (29)$$

where  $\hat{u}_\pm := \frac{1}{2}(1 \pm \hat{\sigma}_3)$  are *mutually annihilating iso-idempotents*. In the standard iso-basis of  $\tilde{\mathbf{G}}_3$ ,

$$\begin{aligned} \{\hat{\sigma}_1, \hat{\sigma}_2 = i\hat{\sigma}_1\hat{\sigma}_3, \hat{\sigma}_3\}, \\ \hat{i} = \hat{\sigma}_1\hat{\sigma}_2\hat{\sigma}_3, \end{aligned} \quad (30)$$




Fig. 1: In the top, we illustrate the structure of the iso-Deuteron as a hadronic bound state of an iso-proton and an iso-neutron in axial triplet coupling, thus representing for the first time the spin of the Deuteron  $S_D = 1$  in its ground state, that with null angular contributions  $L_D = 0$ . The prefix “iso” represents the novel hadronic spin characterized by the iso-Pauli matrices, (14), with an explicit and concrete realization of Bohm’s hidden variable  $\lambda$ . The axial triplet coupling was first identified in the new chemical species of magne-cules (see [49, Chapter 8, page 303 on] and [50, 51]) and then used for the new Intermediate Controlled Nuclear Fusion [52–54]. In the bottom, we illustrate the structure model of the iso-Helium as a hadronic bound state under strong interactions of two iso-Deuterons in singlet coupling, which allows a representation of the null spin and magnetic moment of Helium in its ground state. It should be noted that the above model is not necessarily extendable to heavier stable nuclei due to the prior need of resolving the problem of nuclear stability caused by the natural instability of the neutron, which problem is planned for study in a subsequent paper (see [65] for a preliminary study).

is the *conventional unit* of the associative algebra  $\tilde{\mathbf{G}}_3$ . It must be remembered that  $\hat{\sigma}_k$  verify the property

$$\hat{\sigma}_k^2 = \hat{\sigma}_k \star \hat{\sigma}_k = 1, \tag{31}$$

for  $k = 1, 2, 3$ , where the  $\star$  denotes the iso-product.

We now show that the hidden variable  $\lambda$  of the iso-Pauli matrices (14) can provide a second representation of the deformation of the magnetic moment of nucleons of [57, 58] with consequential exact representation of nuclear magnetic moments.

By introducing the realization of the hidden variable  $\lambda$

$$\lambda = e^\phi \geq 0, \tag{32}$$

with respect to the basis of the *standard unit* of the iso-Pauli matrices  $\hat{I}$ , the *iso-reciprocal*  $\hat{T}$  and the *iso-vector basis*  $\{\hat{\sigma}_k\}$ , are given by

$$\hat{I} = \cosh \phi + \sigma_3 \sinh \phi = e^{\phi \sigma_3}, \tag{33}$$

$$\hat{T} = \cosh \phi - \sigma_3 \sinh \phi = e^{-\phi \sigma_3},$$

Consequently

$$\begin{aligned} \hat{\sigma}_1 &= \sigma_1 \hat{I} = \hat{T} \sigma_1, \\ \hat{\sigma}_2 &= \sigma_2 \hat{I} = \hat{T} \sigma_2, \\ \hat{\sigma}_3 &= \sigma_3 \hat{I} = \hat{I} \sigma_3. \end{aligned} \tag{34}$$

By recalling that  $\sigma_3$  characterizes the nucleon spin  $S = 1/2$ , we reach the result that the replacement of the standard basis of the Clifford algebra  $\mathbf{G}_3$  for Pauli matrices with the iso-Pauli matrices (14) implies the EPR completion of  $\hat{\sigma}_3$  into the expression

$$\hat{\sigma}_3 |\hat{b}\rangle = \sigma_3 \hat{I} |\hat{b}\rangle = \sigma_3 e^{\phi \sigma_3} |\hat{b}\rangle. \tag{35}$$

Recall that the quantum mechanical (qm) relationship between magnetic moments  $\mu$  and spins  $S$  occurs via the gyro-magnetic factor  $g$ ,

$$\mu = gS, \tag{36}$$

and that the corresponding relation for the isotopic branch of hadronic mechanics (hm) is given by an expression of the type [57]

$$\mu_{hm} |\hat{b}\rangle = KgS |\hat{b}\rangle, \tag{37}$$

where  $K$  is an iso-renormalization constant of the gyromagnetic factor  $g$  created by the new notion of hadronic spin  $1/2$ . By using property (28), we reach the relation

$$\mu_{hm} |\hat{b}\rangle = e^{\phi \sigma_3} \mu_{qm} |\hat{b}\rangle = e^{\phi \sigma_3} gS |\hat{b}\rangle. \tag{38}$$

Recall also that: 1) Bohm’s hidden variable  $\lambda$  is associated with the *spin* of a particle according to (14); 2) The proton and the neutron have the same spin  $1/2$  and essentially the same mass, thus being characterized by the same  $\lambda$ ; 3) The quantum mechanical representation of the magnetic moment

of the Deuteron is *in excess* of about 3% according to (24). By selecting the value for conformity with the selected spin orientation (Fig. 1)

$$\sigma_3 |\hat{b}\rangle = -|\hat{b}\rangle, \quad (39)$$

we can write the expression per each nucleon

$$\mu_{hm,k} \approx (1 + \phi\sigma_3)\mu_{qm,k} = (1 - \phi)\mu_{qm,k}, \quad k = p, n, \quad (40)$$

from which we obtain the iso-renormalized value of the magnetic moment of the proton and of the neutron

$$\hat{\mu}_p = +(1 - \phi) 2.79285 \mu_N, \quad \hat{\mu}_n = -(1 - \phi) 1.91304 \mu_N, \quad (41)$$

with corresponding value for the magnetic moment of the Deuteron

$$\begin{aligned} \mu_D^{hm} &= (1 - \phi) 2.79285 - (1 - \phi) 1.91304 \mu_N = \\ &= (1 - \phi) 0.87981 \mu_N = \mu_D^{ex} = 0.85647 \mu_N. \end{aligned} \quad (42)$$

From this, we obtain the numeric value

$$\phi = 1 - 0.87981/0.85647 = 1 - 0.02334 = 0.97666, \quad (43)$$

with corresponding *numeric value of Bohm's hidden variable for the Deuteron*

$$\lambda = e^\phi = e^{0.97666} = 2.65557, \quad (44)$$

by thereby achieving the desired exact representation of the magnetic moment of the Deuteron in terms of Bohm hidden variable  $\lambda$ . Its invariance over time follows from the derivation of iso-Pauli matrices (14) from the Lie-Santilli iso-symmetry  $\hat{\mathcal{P}}(3.1)$  [39]–[41].

The iso-representation of the magnetic moment of  $4 - He - 2$  as the *iso-Helium*  $\widehat{He}(2)$  is a consequence (Fig. 1). The study of the iso-representation for heavier stable nuclei was initiated in [65], but its in-depth achievement requires the still missing consistent representation of nuclear stability against the natural instability of the neutron, which problem is planned for study in a subsequent paper.

We should finally note that in this section we have used the *standard Clifford algebra* and not the full isotopic Clifford algebra  $\hat{\mathbf{G}}$  introduced by R. da Rocha and J. Vaz Jr. [66]. This is due to the fact that the full isotopy  $\hat{\mathbf{G}}_3$  of  $\mathbf{G}_3$  would have required the use of iso-product (6) with the isotopic element  $\hat{T} = e^{-\phi\sigma_3} = 1/\hat{I}$ , and the consequential lack of representation in (38) of the magnetic moment of the Deuteron for spin  $S_D = 1$  in the ground state.

The understanding is however that the full iso-Clifford iso-algebra  $\hat{\mathbf{G}}_{3N}$  is expected to be important for the numerically exact and time invariant representation of the spins and magnetic moments of nuclei with  $A \geq 2$  nucleons.

In a nutshell, we can say that the Copenhagen interpretation of quantum mechanics deals with the simplest possible realization of quantum axioms, while the EPR completion of quantum into hadronic mechanics deals with progressively broader realizations of the same axioms for systems with progressively increasing complexity.

## 4 Acknowledgments

The author would like to thank Prof. G. Sobczyk for consultations in the representation of the isotopic Pauli matrices via the conventional Clifford's algebras. Thanks are also due for penetrating critical comments received from the participants of the *2020 International Teleconference on the EPR argument* and the *2021 International Conference on Applied Category Theory and Graph-Operad-Logic*. Additional thanks are due to various colleagues for technical controls and to Mrs. Sherri Stone for linguistic control of the manuscript. The author is solely responsible for the content of this paper.

Received on May 20, 2022

## References

1. Santilli R.M. Representation of the anomalous magnetic moment of the muons via the Einstein-Podolsky-Rosen completion of quantum into hadronic mechanics. *Progress in Physics*, 2021, v.17, 210–215. <http://www.santilli-foundation.org/muon-anomaly-pp.pdf>.
2. Santilli R.M. Elements of Hadronic Mechanics. Volumes I, Mathematical Foundations. Ukraine Academy of Sciences, Kiev, 1995. <http://www.santilli-foundation.org/docs/Santilli-300.pdf>.
3. Santilli R.M. Elements of Hadronic Mechanics. Volumes II, Theoretical Foundations. Ukraine Academy of Sciences, Kiev, 1994. <http://www.santilli-foundation.org/docs/Santilli-301.pdf>.
4. Santilli R.M. Elements of Hadronic Mechanics. Volumes III, Experimental Verifications. Ukraine Academy of Sciences, Kiev, 2016. <http://www.santilli-foundation.org/docs/elements-hadronic-mechanics-iii.compressed.pdf>.
5. Sourlas D.S. and Tsagas G.T. Mathematical Foundation of the Lie-Santilli Theory. Ukraine Academy of Sciences, Kiev, Ukraine, 1993. <http://www.santilli-foundation.org/docs/santilli-70.pdf>.
6. Ganfornina R.M.F. and Valdes J.N. Fundamentos de la Isotopia de Santilli. International Academic Press, Palm Harbor, FL, 2001. <http://www.i-b-r.org/docs/spanish.pdf>. English translation: Algebras, Groups and Geometries, 2015, v. 32, 135–308 (2015) <http://www.i-b-r.org/docs/Aversa-translation.pdf>.
7. Georgiev S. Foundations of IsoDifferential Calculus. Vol 1. – Iso-Differential and Iso-Integral Calculus for Iso-Functions in One Variable. Vol 2. – Iso-Differential and Iso-Integral Calculus for Iso-Functions in Several Variables. Vol. 3 – Iso-Ordinary Iso-Differential Equations. Vol. 4 – Iso-Differential Equations. Vol.5 – Iso-Stochastic Iso-Differential Equations. Vol. 6 – Theory of Iso-Measurable Iso-Functions. Nova Publishers, New York, NY, 2014 (I), 2014 (II), 2014 (III), 2015 (IV), 2015 (V), 2016 (VI), 2022 (I new ed.).
8. Aringazin A.K., Jannussis A., Lopez F., Nishioka M. and Vel-janosky B. Santilli's Lie-Isotopic Generalization of Galilei and Einstein Relativities. Kostakaris Publishers, Athens, Greece, 1991. <http://www.santilli-foundation.org/docs/Santilli-108.pdf>.
9. Kadeisvili J. V. Santilli's Isotopies of Contemporary Algebras, Geometries and Relativities, Second edition. Ukraine Academy of Sciences, Kiev, 1997. <http://www.santilli-foundation.org/docs/Santilli-60.pdf>.
10. Gandzha I. and Kadeisvili J. V. New Sciences for a New Era: Mathematical, Physical and Chemical Discoveries of Ruggero Maria Santilli. Sankata Printing Press, Nepal, 2011. <http://www.santilli-foundation.org/docs/RMS.pdf>.
11. Santilli R. M. Studies on A. Einstein, B. Podolsky, and N. Rosen prediction that quantum mechanics is not a complete theory. I: Basic methods. *Ratio Mathematica*, 2020, v.38, 5–69. <http://www.eprdebates.org/docs/epr-review-i.pdf>.

12. Santilli R. M. Studies on A. Einstein, B. Podolsky, and N. Rosen prediction that quantum mechanics is not a complete theory. II: Apparent proof of the EPR argument. *Ratio Mathematica*, 2020, v. 38, 71–138. <http://www.eprdebates.org/docs/epr-review-ii.pdf>.
13. Santilli R. M. Studies on A. Einstein, B. Podolsky, and N. Rosen prediction that quantum mechanics is not a complete theory. III: Illustrative examples and applications. *Ratio Mathematica*, 2020, v. 38, 139–222. <http://www.eprdebates.org/docs/epr-review-iii.pdf>.
14. Santilli R. M. Isorepresentation of the Lie-isotopic SU(2) Algebra with Application to Nuclear Physics and Local Realism. *Acta Applicandae Mathematicae*, 1998, v. 50, 177–190. <http://www.santilli-foundation.org/docs/Santilli-27.pdf>.
15. Santilli R. M. Studies on the classical determinism predicted by A. Einstein, B. Podolsky and N. Rosen. *Ratio Mathematica*, 2019, v. 37, 5–23. <http://www.eprdebates.org/docs/epr-paper-ii.pdf>.
16. Einstein A., Podolsky B., and Rosen N. Can quantum-mechanical description of physical reality be considered complete? *Phys. Rev.*, 1935, v. 47, 777–791. <http://www.eprdebates.org/docs/epr-argument.pdf>.
17. Beghella-Bartoli S. and Santilli R. M., eds, Proceedings of the 2020 Teleconference on the Einstein-Podolsky-Rosen argument that “Quantum mechanics is not a complete theory”. Curran Associates, New York, NY, 2021. <http://www.proceedings.com/59404.html> (printed), 60007.html (electronic). <http://www.santilli-foundation.org/epr-table-of-content.pdf> (TOC).
18. Santilli R. M. Overview of historical and recent verifications of the EPR argument and their applications to physics, chemistry and biology. APAV - Accademia Piceno Aprutina dei Velati, Pescara, Italy, 2021. <http://www.santilli-foundation.org/epr-overview-2021.pdf>.
19. Dunning-Davies J. A Present Day Perspective on Einstein-Podolsky-Rosen and its Consequences. *Journal of Modern Physics*, 2021, v. 12, 887–936.
20. Santilli R. M. Apparent Unsettled Value of the Recently Measured Muon Magnetic Moment. *Progress in Physics*, 2022, v. 18, 15–18. <http://www.santilli-foundation.org/docs/muon-meanlife-2022.pdf>
21. Aronson S. H., Bock G. J., Cheng J. Y., and Fischbach E. Energy dependence of the fundamental parameters of the  $K^0 - \bar{K}^0$  system. *Phys. Rev. D*, 1983, v. 28, 495–503.
22. Santilli R. M. Embedding of Lie-algebras into Lie-admissible algebras. *Nuovo Cimento*, 1967, v. 51, 570. <http://www.santilli-foundation.org/docs/Santilli-54.pdf>.
23. Blokhintsev D. I. The Philosophy of Quantum Mechanics. JINR (in Russian), 1965. Springer (in English), 1968.
24. Bohm D. A Suggested Interpretation of the Quantum Theory in Terms of “Hidden Variables”. *Physical Review*, 1952, v. 85, 166–182. <http://journals.aps.org/pr/abstract/10.1103/PhysRev.85.166>.
25. Blatt J. M. and Weisskopf V. F. Theoretical Nuclear Physics. Wiley and Sons, 1952.
26. Santilli R. M. On a possible Lie-admissible covering of Galilei’s relativity in Newtonian mechanics for nonconservative and Galilei form-non-invariant systems. *Hadronic J.*, 1978, v. 1, 223–423. <http://www.santilli-foundation.org/docs/Santilli-58.pdf>.
27. Santilli R. M. Need of subjecting to an experimental verification the validity within a hadron of Einstein special relativity and Pauli exclusion principle. *Hadronic J.*, 1978, v. 1, 574–901. <http://www.santilli-foundation.org/docs/santilli-73.pdf>.
28. Santilli R. M. Foundation of Theoretical Mechanics. Vol. I, The Inverse Problem in Newtonian Mechanics. Springer-Verlag, Heidelberg, Germany, 1978. <http://www.santilli-foundation.org/docs/Santilli-209.pdf>.
29. Santilli R. M. Foundation of Theoretical Mechanics. Vol. II, Birkhoffian Generalization of Hamiltonian Mechanics. Springer-Verlag, Heidelberg, Germany, 1983. <http://www.santilli-foundation.org/docs/santilli-69.pdf>.
30. Kadeisvili J. V. An introduction to the Lie-Santilli isotopic theory. *Mathematical Methods in Applied Sciences*, 1996, v. 19, 134–168. <http://www.santilli-foundation.org/docs/Santilli-30.pdf>.
31. Mukhtibodh A. S. and Santilli R. M. Studies of the Regular and Irregular Isorepresentations of the Lie-Santilli Isotheory. *Journal of Generalized Lie Theories*, 2007, v. 11, 1–32. <http://www.santilli-foundation.org/docs/isorep-Lie-Santilli-2017.pdf>.
32. Santilli R. M. Lie-isotopic Lifting of Special Relativity for Extended Deformable Particles. *Lettere Nuovo Cimento*, 1983, v. 37, 545–555. <http://www.santilli-foundation.org/docs/Santilli-50.pdf>.
33. Santilli R. M. Lie-isotopic Lifting of Unitary Symmetries and of Wigner’s Theorem for Extended and Deformable Particles. *Lettere Nuovo Cimento*, 1983, v. 38, 509–516. <http://www.santilli-foundation.org/docs/Santilli-51.pdf>.
34. Santilli R. M. Isotopies of Lie Symmetries. I: Basic theory. *Hadronic J.*, 1985, v. 8, 8–33. <http://www.santilli-foundation.org/docs/santilli-65.pdf>.
35. Santilli R. M. Isotopies of Lie Symmetries. II: Isotopies of the rotational symmetry. *Hadronic J.*, 1985, v. 8, 36–59. <http://www.santilli-foundation.org/docs/santilli-65.pdf>.
36. Santilli R. M. Rotational isotopic symmetries. ICTP Communication No. IC/91/261, 1991. <http://www.santilli-foundation.org/docs/Santilli-148.pdf>.
37. Santilli R. M. Isotopic Lifting of the SU(2) Symmetry with Applications to Nuclear Physics. *JINR Rapid Comm.*, 1993, v. 6, 24–36. <http://www.santilli-foundation.org/docs/Santilli-19.pdf>.
38. Santilli R. M. Isotopic generalization of the Poincaré symmetry. ICTP Communication No. IC/91/45, 1991. <http://www.santilli-foundation.org/docs/Santilli-140.pdf>.
39. Santilli R. M. Nonlinear, Nonlocal and Noncanonical Isotopies of the Poincaré Symmetry. *Moscow Phys. Soc.*, 1993, v. 3, 255–289. <http://www.santilli-foundation.org/docs/Santilli-40.pdf>.
40. Santilli R. M. Isotopies of the spinorial covering of the Poincaré symmetry. Communication of the Joint Institute for Nuclear Research, Dubna, Russia, No. E4-93-252, 1993.
41. Santilli R. M. Relativistic synthesis of the neutron from the hydrogen atom via the isospinorial covering of the iso-Poincaré symmetry. *Chinese J. System Eng. and Electr.*, 1995, v. 6, 177–202. <http://www.santilli-foundation.org/docs/Santilli-18.pdf>.
42. Santilli R. M. Iso-Minkowskian Geometry for the Gravitational Treatment of Matter and its Isodual for Antimatter. *Intern. J. Modern Phys. D*, 1998, v. 7, 351–385. <http://www.santilli-foundation.org/docs/Santilli-35.pdf>.
43. Bohm D. A Suggested Interpretation of the Quantum Theory in Terms of “Hidden Variables”. *Physical Review*, 1952, v. 85, 166–182. [journals.aps.org/pr/abstract/10.1103/PhysRev.85.166](http://journals.aps.org/pr/abstract/10.1103/PhysRev.85.166).
44. Bell J. S. On the Einstein Podolsky Rosen paradox. *Physics*, 1964, v. 1, 195–221.
45. Myung H. C. and Santilli R. M. Modular-isotopic Hilbert space formulation of the exterior strong problem. *Hadronic Journal*, 1982, v. 5, 1277–1366. <http://www.santilli-foundation.org/docs/myung-santilli-1982.pdf>.
46. Santilli R. M. Isonumbers and Genonumbers of Dimensions 1, 2, 4, 8, their Isoduals and Pseudoduals, and “Hidden Numbers” of Dimension 3, 5, 6, 7. *Algebras, Groups and Geometries*, 1993, v. 10, 273–322. <http://www.santilli-foundation.org/docs/Santilli-34.pdf>.
47. Santilli R. M. Invariant Lie-isotopic and Lie-admissible formulation of quantum deformations. *Found. Phys.*, 1997, v. 27, 1159–1177. <http://www.santilli-foundation.org/docs/Santilli-06.pdf>.
48. Goldstein S. Bohmian (de Broglie-Bohm) Mechanics. Stanford Encyclopedia of Philosophy, 2021. <http://plato.stanford.edu/entries/qm-bohm/>

49. Santilli R. M. Foundations of Hadronic Chemistry, with Applications to New Clean Energies and Fuels. Kluwer Academic Publishers, 2001. <http://www.santilli-foundation.org/docs/Santilli-113.pdf>. Russian translation: Aringazin A. K. [i-b-r.org/docs/Santilli-Hadronic-Chemistry.pdf](http://i-b-r.org/docs/Santilli-Hadronic-Chemistry.pdf).
50. Pandhurneka C. P. and Zodape S. P. Santilli's Magneucles and Their Applications. *American Journal of Modern Physics*, 2017, v. 6, 64–73. <http://www.santilli-foundation.org/docs/magneucles-2017.pdf>.
51. Yang Y., Kadeisvili J. V. and Marton S. Experimental Confirmations of the New Chemical Species of Santilli Magneucles. *The Open Physical Chemistry Journal*, 2013, v. 5, 1–15. <http://www.santilli-foundation.org/docs/Magneucles-2012.pdf>.
52. Santilli R. M. The novel Intermediate Controlled Nuclear fusions, a report for its industrial realization. *Hadronic Journal*, 2008, v. 31, 15–43. <http://www.santilli-foundation.org/docs/CNF-HJ.pdf>.
53. Brenna R., Kuliczowski T. and Ying L. Verification of Santilli intermediate Controlled Nuclear Fusions without harmful radiations and the production of magneucular clusters. *New Advances in Physics*, 2011, v. 5, 9–21. <http://www.santilli-foundation.org/docs/ICNF-2.pdf>.
54. Lynch C. and Yang Y. Confirmations of Santilli Intermediate Nuclear Fusions of Deuteron and Carbon into Nitrogen without Radiations. *The Open Physical Chemistry Journal*, 2013, v. 5, 17–39. <http://www.santilli-foundation.org/docs/ICNF-Conf-2013.pdf>.
55. Vonsovsk S. Magnetism of Elementary Particles. Mir Publishers, 1975.
56. Fermi E. Nuclear Physics. University of Chicago Press, 1949.
57. Santilli R. M. A quantitative isotopic representation of the deuteron magnetic moment. in Proceedings of the International Symposium "Dubna Deuteron-93". Joint Institute for Nuclear Research, Dubna, Russia, 1994. <http://www.santilli-foundation.org/docs/Santilli-134.pdf>.
58. Santilli R. M. The Physics of New Clean Energies and Fuels According to Hadronic Mechanics. *Journal of New Energy, Special Issue*, 1998, 318 pages. <http://www.santilli-foundation.org/docs/Santilli-114.pdf>.
59. Rauch H. and Zeilinger A. Demonstration of SU(2) symmetry by neutron interferometers. *Hadronic Journal*, 1981, v. 4, 1280–1285. <http://www.santilli-foundation.org/docs/rauch-zeilinger-1981.pdf>.
60. Hestenes D. and Sobczyk G. Clifford Algebra to Geometric Calculus: A Unified Language for Mathematics and Physics, 2nd ed. Kluwer, 1992.
61. Doran C. and Lasenby A. Geometric Algebra for Physicists. Cambridge University Press, 2003.
62. Sobczyk G. New Foundations in Mathematics: The Geometric Concept of numbers. Birkhauser, New York, 2013.
63. Sobczyk G. Spacetime Vector Analysis. *Physics Letters A*, 1981, v. 84, 45–49.
64. Sobczyk G. Conformal Mappings in Geometric Algebra. *Notices of the AMS*, 2012, v. 59, 264–273.
65. Bhalekar A. A. and Santilli R. M. Exact and Invariant representation of nuclear magnetic moments and spins according to hadronic mechanics. *American Journal of Modern Physics*, 2016, v. 5, 56–92. <http://www.santilli-foundation.org/docs/nuclear-MM-spins.pdf>.
66. da Rocha R. and Vaz J. Jr. Isotopic liftings of Clifford algebras and applications in elementary particle mass matrices. *Int. J. Theor. Phys.*, 2007, v. 46, 2464–2487. arXiv: math-ph/0710.0832.